

Factsheet on environmental exposure and offspring health

Published March 2024

Key facts

- The health of the offspring can be affected by environmental stressors through maternal exposure in the preconception period, prenatal and neonatal life, but also through paternal exposure in the preconception period. Environmental exposure includes exposure at the workplace.
- Exposure to environmental stressors during early childhood can have immediate and future health consequences.
- Reducing and potentially eliminating the exposure of children to environmental stressors in the early years of their lives, and even before conception and pregnancy, can protect their health and that of future generations.

Background information

Throughout their lifetimes, humans are exposed to a multitude of human-made chemicals. Increasing evidence suggests that exposure to environmental and occupational factors, not only during pregnancy but also prior to conception, can influence the health of offspring [1]. Research indicates that exposures can impact germ cells and might even induce potentially heritable alterations of the epigenome, in both males and females, potentially leading to adverse effects on the health and development of future generations. Since it is well-established that disruptions during early development can have consequences that manifest later in life, it is imperative to underscore the significance of addressing environmental exposures to safeguard the health and well-being of offspring.

Of note, many chemicals have not been appropriately or never been tested for toxicity to the developing foetus.

Effects of environmental exposure on the general health of offspring

During pregnancy, the impact on the offspring hinges not only on the nature and duration of exposure to environmental stressors, but also on the stage of foetal development. Additionally, concerns arise regarding mother's exposure to bioaccumulative chemicals, which have the potential to accumulate in fat tissue, liver, or lungs over extended periods [2]. These chemicals

may be released during breastfeeding when fat tissue is mobilised, raising further health concerns.




Figure 1: Potential effects of exposure to environmental factors [1,2]; EDCs, Endocrine Disrupting Chemicals (Created with BioRender.com).

→ **Parental exposure to pesticides and air pollution are linked with childhood leukaemia.**

- Maternal and paternal exposure to pesticides before and/or during pregnancy as well as air pollution have been identified as factors linked to an increased risk of acute childhood leukaemia.

→ **Exposure to chemicals is linked with impaired neurobehavioral development.**

- Prenatal exposure to chemicals such as endocrine disrupting chemicals (EDCs), alcohol, and organophosphate pesticides has been linked to compromised cognitive development [3]. Furthermore, exposure to other toxic agents has been linked to increased risks of attention deficit/hyperactivity disorders and/or autism spectrum disorders in the developing offspring.

→ **Pesticide exposure may lead to metabolic syndrome/obesity in the offspring.**

- Prenatal exposure to certain substances, including EDCs, pesticides and insecticides, can elevate the risk of obesity and metabolic syndrome in offspring by affecting their energy metabolism.

→ **Maternal occupational exposure is associated with congenital malformations.**

- Maternal occupational exposure to solvents, occurring from three months before conception until the end of the first trimester, is associated with an increased risk of neural tube defects in offspring.

→ **Persistent organic pollutants alter the immune system.**

- Early-life exposure, both pre- and postnatal, to persistent organic pollutants has the potential to disrupt the developing immune and respiratory systems, reducing the offspring's capacity to combat infections and heightening the risk of allergic manifestations such as asthma.

Effect of environmental exposure on the reproductive health of offspring

→ **Male offspring: effect on genital anomalies, testicular function and testis cancer.**

- Exposure in early gestation to factors such as maternal smoking, maternal stress, chemicals with anti-androgenic properties or mild analgesics have been implicated in disrupting the normal differentiation and growth of male reproductive organs.
- Prenatal exposure to EDCs has been associated with altered testicular descent, leading to cryptorchidism. Substances like glycol ethers can directly interfere with urethra or penis formation, resulting in conditions such as hypospadias or micropenis. Bisphenol A (BPA) has been linked to impaired semen quality and testicular function in young adult men, as well as with an increased incidence of testicular diseases and testicular germ cell tumours.

→ **Female offspring: effect on ovarian function and genital anomalies with a transgenerational effect.**

- Numerous studies have connected gestational exposure to environmental and occupational stressors, including EDCs, air pollution, and cigarette smoking, with postnatal health issues in women. Exposure to substances like diethylstilbesterol (DES), phthalates, and bisphenols has been linked to earlier puberty onset, an increased risk of polycystic ovary syndrome and endometriosis, disrupted menstrual cycles, and infertility or longer time-to-pregnancy in female offspring.
- Furthermore, adult uterine abnormalities can be influenced by developmental effects induced by EDCs, possibly mediated by changes in genes relevant to uterine structure and endometrial function, with potential transgenerational implications [4, 5].

Time for action: policies to reduce the effect of environmental exposure on offspring health.

Facilitate Research

- ✓ Foster research to uncover correlations between environmental and occupational sources of exposure, such as air pollution, and health outcomes in children.
- ✓ Improve surveillance by establishing channels for reporting health-related concerns and suspected cases of disease caused by environmental exposure.

Promote Awareness

- ✓ Disseminate knowledge and educate families and communities about the most common environmental risks and measures to prevent them.

Support Prevention

- ✓ Develop and enforce prevention strategies to mitigate health risks linked to environmental factors like air pollution and hazardous chemicals.
- ✓ Increase the requirements for evaluating the developmental toxicity and subsequent approval of chemicals to be used by consumers and workers.
- ✓ Stay abreast of emerging environmental research and concerns to inform policy decisions.

References¹

1. Schmitz-Felten, E., et al., *State of the art report on reproductive toxicants*. European Risk Observatory Summary. Luxembourg: European Agency for Safety and Health at Work, 2016: p. 1-61.
2. Vinnars, M.T., M. Bixo, and P. Damdimopoulou, *Pregnancy-related maternal physiological adaptations and fetal chemical exposure*. *Mol Cell Endocrinol*, 2023. **578**: p. 112064.
3. Caporale, N., et al., *From cohorts to molecules: Adverse impacts of endocrine disrupting mixtures*. *Science*, 2022. **375**(6582): p. eabe8244.
4. Miranda, M.L., et al., *Blood lead levels among pregnant women: historical versus contemporaneous exposures*. *Int J Environ Res Public Health*, 2010. **7**(4): p. 1508-19.
5. Dong, Y., et al., *Exposure patterns, chemical structural signatures, and health risks of pesticides in breast milk: A multicenter study in China*. *Sci Total Environ*, 2022. **830**: p. 154617.

¹Only key papers were included. Further detailed references on individual studies can be requested from ESHRE by contacting guidelines@eshre.eu

Developed by the ESHRE expert panel for environmental factors and fertility, consisting of Willem Ombelet, Maribel Acién, Pauliina Damdimopoulou, Linda Giudice, Niels E. Skakkebaek, Miguel Angel Checa, Rémi Béranger, Nicolas Garrido, Karin Sørig Hougaard, Rune Lindahl-Jacobsen, Olwenn Martin, Jeanne Perrin, Cristina Richie, and Nathalie Vermeulen.